

Colloquium

University of Notre Dame
Department of Mathematics

Speaker: Marta Lewicka

University of Pittsburgh

Will give a lecture entitled

Dimension reduction via Gamma-convergence of non-convex energies, and the quantitative immersability of Riemannian metrics.

Date: Thursday, January 16, 2020

Time: 4:00 PM

Location: 127 Hayes-Healy Hall

Departmental Tea: Tea in Room 257 (lounge in Hurley Hall) at 3:30 p.m.



Abstract:

In this talk, we present results that relate the following three contexts: (i) Given a Riemannian metric G we study the problem of finding the infimum of the averaged pointwise deficit of an immersion from being an orientation-preserving (equi-dimensional) isometric immersion of G . We are also interested in discovering the structure and regularity of the optimal immersion. Questions of this type can be seen as a quantitative variation of the very classical topics in Differential Geometry: existence, rigidity and flexibility of isometric immersions. (ii) Given a family of non-convex energies E^h , parametrized by a three-dimensional plate's thickness h , we are interested in finding the scaling laws of the infima of E^h , in terms of the powers of h , as h goes to 0. Subsequently, we want to derive the Gamma-limits (variational limits) of E^h and reveal the asymptotic structure of the minimizing sequences. This topic is classical in the Calculus of Variations and may also be seen as a variation of the study of nonlinear elasticity. Combining (i) and (ii), we first show how the deficit mentioned in (i) can be measured by the energies E^h in (ii), and perform the full scaling analysis of E^h , in the context of dimension reduction. Then, we derive the Gamma-limits of $h^{-2n}E^h$ for all powers n . We show the energy quantization, in the sense that the even powers $2n$ of h are the only possible ones (all of them are also attained). For each n , we identify conditions for the validity of the corresponding scaling, in terms of the vanishing of Riemann curvatures of G up to appropriate orders, and in terms of the singular behavior of the minimizing immersions. If time permits, we will explain how the above problems are linked to: (iii) The description of elastic materials exhibiting residual stress at free equilibria. Examples of such structures and their actuations include: plastically strained sheets; specifically engineered swelling or shrinking gels; growing tissues, and atomically thin graphene layers. Our results and methods display the interaction of nonlinear pdes, geometry and mechanics of materials in the prediction of patterns and shape formation.